

Postoperative Instructions for Open Nasal Reconstruction

- 1. This procedure is performed to attempt correction of an external and internal nasal deformity associated with nasal obstruction. These procedures are complex and may require 3-4 hours of general anesthesia.
- 2. Bloody nasal drainage may last for a few days. Change the nasal drip pad as needed.
- 3. There will be a cast over the nose. There will be splints sutured in the nose. These will be removed 1-2 weeks after surgery. The nose will be very congested. There is nothing that may be done to resolve this quickly. It will improve over the weeks after surgery.
- 4. There will be sutures between the nostrils. These sutures are removed 1 week after surgery. Clean the incision and sutures 3 times a day with peroxide and apply over the counter Bacitracin antibiotic ointment.
- 5. No nose blowing for 2 weeks. No strenuous activity for 2 weeks. No contact sports for 6 weeks.
- 6. Expect some bruising over the nose and around the eyes. There will be external nasal swelling. The most noticeable swelling will improve over 2-3 weeks. However, the nose may not reach final appearance for 6 months.
- 7. For the first 48 hours after surgery, apply an ice pack over the nose for 20 minutes every 2 hours.
- 8. Sleep with your head elevated in a recliner or on 3-4 pillows for 1 week.
- 9. Pain medications and antibiotics will be prescribed. You may also take Advil or Motrin.
- 10. Fevers for a few days after surgery may occur. Use Advil or Motrin.
- 11. The upper midline teeth may be numb for a few weeks after surgery.
- 12. Nasal obstruction may be multifactorial. Other procedures may be required to further improve nasal breathing problems after this surgery.

I provided before the date of surgery.	, confirm that I have read and understand the information
Signature	 Date