



CAPITAL CITY
PERIODONTICS & ORAL IMPLANTOLOGY

*** For your surgical appointment we recommend that you park on the **5th Floor of the Huntington Center Parking Garage** in one of our patient reserved parking spots which are numbered **501, 502 or 503**. The reserved parking spots for CCP surgical patients have **YELLOW RESERVED** signs. Please look for the spots to be on the **RIGHT** as you are driving through the garage.***

The more prepared a dental patient is prior to therapy and having a realistic idea of what to expect after treatment will help the person to be more relaxed for their procedure. It is important to follow the dentist's postoperative instructions to allow for your healing to go as quickly and smoothly as possible.

All treatment appointments are chosen based on each patient's specific dental needs. Please be aware that we kindly request a 48 business hour notice for any surgical appointment change. A fee may be assessed if we do not receive sufficient notice of change.

Pre-treatment checklist of recommendations for all surgical procedures

- NO smoking or tobacco usage for 48 hours prior to treatment; tobacco negatively affects the level of comfort during the appointment.
- NO alcohol 48 hours prior to treatment; the dentist cannot safely treat you if you have had alcohol, narcotic pain medication, or any street or recreational drugs.
- NO caffeine 24 hours prior.
- Review and take ALL prescriptions medications (especially blood pressure medications) as directed. Unless you are given other specific instructions.
- Avoid any over the counter medications which may contribute to blood thinning for **one week prior** to your procedure, unless otherwise instructed including, but not limited to: Advil, Aleve, Aspirin, Celebrix, Feldene, Herbal Supplements, Ibuprofen, Motrin, Multi-Vitamins, and Nuprin.

Pre-treatment checklist if IV Sedation is being performed with your surgical procedure

- No eating or drinking for 6 hours prior to the procedure. Take **all** prescription medications with a small amount of water (**especially blood pressure medications**).
- Notify us prior to treatment of **all** allergies including but not limited to: Fentanyl, Versed, or Phenergan.
- All diabetics taking Insulin, should take ½ the dosage of their Insulin in the morning of the dental appointment. Diabetic patients taking Insulin will have early schedule priority. Sugar level should be checked the morning of the appointment.
- Remember to remove any fingernail polish or unnatural nail covering.



CAPITAL CITY
PERIODONTICS & ORAL IMPLANTOLOGY

- Please leave jewelry and watches at home.
- When possible please do not wear contact lenses.
- Wear short sleeves and comfortable clothing. Bring a light weight button-up or zipper jacket or sweater. Wear comfortable shoes and socks. Please do not wear white or expensive clothing to your surgical appointment.
- It is essential that you have an escort who drives you to & from your appointment. **You will NOT be able to drive for the remainder of the day.** A blanket and pillow in the car is advisable for comfort on your return trip home.
- For your safety, you must have someone with you at home for the remainder of the day to monitor you.**

Post LANAP Therapy Homecare Recommendations

We have scheduled a significant block of time for our team to concentrate on your visit with us. Please understand that this is surgery, done in a way with laser technology. Healing occurs at an individual's own rate and is affected by outside influences such as mechanical or bacterial trauma. It is very important not to disturb the fibrin clot that seals the pocket and acts as a bandage to allow the bone and tissue to regenerate properly. So please follow the home care and diet instructions carefully.

1. Avoid using straws and spitting for 14 days. The sucking action will cause a vacuum in the mouth and may dislodge the body's natural clotting process.
2. No vigorous exercising, heavy lifting, or any other activities that could cause your heart rate to become elevated for 1 week. These types of activities will increase your blood pressure and possibly dislodge the fibrin clots which are essential to healing. Wait 1 week before resuming or engaging in heavy athletics.
3. No usage of any tobacco / nicotine products for 48 hours post treatment. This includes the use of e-cigarettes and vapor products.
4. Some oozing of blood may occur and will appear to be greatly exaggerated when it dissolves in the saliva. This will almost always stop without attention. If it continues, locate the area involved and apply firm pressure to this area with a gauze or moist teabag for 30 minutes.
5. Do not be alarmed if beginning two weeks after therapy and extending as long as one year or more, the teeth may be sore, tender or sensitive as bone and ligaments around the teeth regenerate and become more firm. This is a sign of healing, but may indicate a bite imbalance that needs to be adjusted. Occlusal adjustments are an important part of your care and maintenance.
6. An oral rinse has been provided; please rinse for 7 – 10 days after your surgery. Gently swish in the morning and the evening for 60 seconds, *NO EATING OR DRINKING* for 20 minutes after rinsing.
7. Oral Hygiene: **Do not use an electric toothbrush until notified by the doctor.** No flossing the surgery site(s) for 1 month.



Post Treatment Medication

1. Some soreness, tenderness, or tooth sensitivity is normal. The discomfort is usually mild to moderate and much less than with conventional blade and suture surgery. To reduce inflammation and promote healing, you will normally be advised to take Ibuprofen 600mg or Tylenol 1000mg four times per day for two days unless you are allergic to Aspirin or Ibuprofen. If Ibuprofen is not sufficient then Tylenol 1000mg can be taken at the same time with the Ibuprofen.
2. Prescriptions may be dispensed prior to treatment for your convenience. Please do not start medications until **after** treatment unless notified otherwise.
3. Please take the medication as prescribed following the directions as your pharmacist has recommended.
4. Consume adequate water while taking medications. This will help prevent nausea, an upset stomach, and expedite the medication's effects.
5. Post-treatment medications may include: Norco, Percocet, Amoxicillin, Clindamycin or Zithromax. **IF YOU ARE ALLERGIC TO ANY OF THESE MEDS INFORM US IMMEDIATELY.**
6. If an antibiotic is prescribed at any time during treatment, you should take the medication exactly as directed until prescription completion.
7. Limit alcohol consumption and sun exposure as they may interfere with the medications prescribed.
8. Do **not** exceed the recommended dosage. Take only the medication as prescribed on the prescription.
9. Do **not** give any of your medication to ANY other individual.
10. Female patients that are prescribed antibiotics are recommended to seek alternative birth control methods the entire menstrual cycle if taking contraceptives.
11. Notify the doctor if you have symptoms which may be related to an allergic reaction such as: skin rash, itching or hives, elevated temperature, increased and/or erratic heart rate, nausea or vomiting, dizziness or fainting, or blurred vision.

Dietary Recommendations after LANAP Treatment

1. First Day: liquid like diet. Use a blender as needed. **DO NOT DRINK THROUGH A STRAW**
2. Next 6 Days: Use soft food suggestions listed below
 - Anything put through a food blender
 - Cream of wheat, oatmeal, grits
 - Mashed potatoes or baked potatoes (ok with butter/ sour cream)
 - Mashed banana, avocado, applesauce or any mashed or blended fruit (except berried fruits)
 - Broth or creamed soups



CAPITAL CITY
PERIODONTICS & ORAL IMPLANTOLOGY

- Mashed or steamed vegetables
 - Mashed yams, baked sweet potato or butternut squash
 - Creamy peanut butter (without solid pieces)
 - Cottage cheese, cream or soft cheeses
 - Eggs any style
 - Yogurt, jell-o, pudding, ice cream, milk shakes or smoothies
 - Ensure, slim fast or any nutritional drinks
3. After 7 Days: Gradually add back softer foods like pasta, chicken and fish
 4. After 14 Days: Resume more normal diet
 5. For 30 Days: DO NOT...chew gum, candy, cookies, chips, popcorn, nuts, anything hard or crunchy, anything that has seeds or hard pieces, meat that shreds and can lodge under the gum and between teeth, raw vegetables or salad.

Notify the office if any of the following occurs:

- There is a constant flow of blood that cannot be stopped after applying pressure for 45 minutes.
- Pain does not decrease after 72 hours.
- Swelling increases after the third day or is significant when looking in the mirror.
- You notice that there is a bad odor or taste in your mouth.
- You experience a fever after 72 hours.

In the event that you are experiencing uncontrollable bleeding or suspect that you have an infection contact us immediately!

During the day please call the office: **614-591-0667**

In the case of a dental emergency after hours please also call the main office phone number & leave a message. When you leave a message we will receive a notification with your voicemail and will return your call based on the nature of your call.

Go to the emergency room if you are unsuccessful in reaching the office after hours.

If you are experiencing a life threatening emergency, please dial 911 immediately.