

ANTERIOR CRUCIATE LIGAMENT AND POSTERIOR CRUCIATE LIGAMENT COMBINED RECONSTRUCTION SURGERY REHABILITATION PROGRAM

I. IMMEDIATE POSTOPERATIVE PHASE (Day 1 - 13)

Goals:

- Restore full passive knee extension
- Diminish joint swelling and pain
- Restore patellar mobility
- Gradually improve knee flexion
- Re-establish quadriceps control
- Restore independent ambulation

POSTOPERATIVE DAY 1 - 4

- Brace:
 - Drop lock brace locked at zero degrees extension with compression wrap
 - Sleep in brace
- Weightbearing:
 - Two crutches as tolerated (less than 50%)
- Range of Motion:
 - Range of motion 0 – 45/50°
 - CPM 0 - 60° as tolerated
- Exercises:
 - Ankle pumps
 - Quad sets
 - Straight leg raising – flexion, abduction, adduction in brace
 - Muscle stimulation to quadriceps (4 hours per day) during quad sets & straight leg raises
 - Patellar mobilizations 5-6x daily
- Ice and elevation every 20 minutes of each hour with knee in extension

POSTOPERATIVE DAY 5 – 13

- Brace:
 - Continue use of drop lock brace locked at zero degrees extension during ambulation & sleep
- Weightbearing:
 - Two crutches: gradually increase WB to 50% by day 7; 75% by day 12
- Range of motion:
 - Day 5: 0 - 65°
 - Day 7: 0 - 75°
 - Day 10: 0 - 85/90°
 - Day 13: 0 - 90°
 - Gradually increase CPM ROM 0 - 70° day 7; 0 - 90° day 12
- Exercises:

- Continue previous exercises
 - Initiate knee extension 60 - 0°
 - Continue use of muscle stimulation
 - Patellar mobilizations 5-6x daily
-
- Continue use of ice, elevation, and compression

II. MAXIMUM PROTECTION PHASE (Week 2 to 6)

Criteria to Enter Phase II:

1. Good quad control (ability to perform good quad set and SLR)
2. Full passive knee extension
3. PROM 0-90 degrees
4. Good patellar mobility
5. Minimal joint effusion

Goals: Control deleterious forces to protect grafts
Nourish articular cartilage
Decrease swelling
Decreased fibrosis
Prevent quad atrophy
Initiate proprioceptive exercises

WEEK 2

- Brace:
 - Continue use of brace locked at zero degrees of extension
 - Weightbearing:
 - As tolerated; approximately 75% body weight
 - Range of Motion:
 - Continue to perform passive ROM 5-6x daily
 - Day 14: 0 - 90°
 - Exercises:
 - Continue quad sets & straight leg raises
 - Continue knee extension 60 - 0°
 - Multi-angle isometrics at 80°, 60° and 40°
 - Patellar mobilizations 5-6x daily
 - Well leg bicycle
 - Weight shifts
 - Mini-squats (0 - 45°)
 - Continue use of muscle stimulation
-
- Continue ice, elevation, and compression

WEEK 3

- Continue above mentioned exercises
- ROM: 0 - 90°
- Continue use of 2 crutches - 75-80% body weight

WEEK 4

- Brace:
 - Continue use of brace locked at zero degrees extension
 - Discontinue sleeping in brace
- Weightbearing:
 - Progress to weight bearing as tolerated with 1 crutch
- Range of Motion:
 - AAROM, PROM: 0 – 90/100°
- Exercises:
 - Weight shifts
 - Mini-squats (0 - 45°)
 - Knee extension 90 - 40° (therapist discretion)
 - Light pool exercises and walking
 - Initiate bicycle for ROM & endurance
 - Begin leg press 60 - 0° (light weight)
 - Proprioception/balance drills
- KT-2000 testing performed – 20 pound (at 25° and 70°)

WEEK 5 – 6

- Discontinue use of crutches week 5 – 6
- Unlock brace for ambulation week 6
- Fit for functional ACL/PCL brace
- Range of Motion week 5: 0 - 105°; week 6: 0 - 115°
- Continue pool exercises
- Initiate lateral lunges
- Hip abduction and adduction
- KT-2000 testing performed – 30 pounds at week 6

III. MODERATE PROTECTION PHASE (Week 7 – 12)

Criteria to Enter Phase III:

1. PROM 0 – 115 degrees
2. Full weightbearing
3. Quadriceps strength > 60% contralateral side (isometric test at 60°)
4. Unchanged KT test (+1 or less)
5. Minimal to no full joint effusion

6. No joint line or patellofemoral pain

Goals: Control forces during ambulation
Progress knee range of motion
Improve lower extremity strength
Enhance proprioception, balance, and neuromuscular control
Improve muscular endurance
Restore limb confidence and function

- Brace:
 - Continue use of unlocked brace for ambulation – discharge week 7 – 8
- Range of Motion:
 - AAROM/PROM 0 - 125°
- Exercises:
 - Continue previous exercises
 - Initiate swimming
 - Initiate lateral and front step-ups (2" step, gradually increase)
 - Progress closed kinetic chain exercises (squats 0 - 60°, leg press 90 - 0°)
 - May begin light hamstring isotonics week 8
 - Progress proprioceptive training

- KT-2000 test: 20 & 30 pounds at week 6 & 8

IV. CONTROLLED ACTIVITY PHASE (Week 13 – 16)

Criteria to Enter Phase IV:

1. AROM 0 - 125°
2. Quadriceps strength > 60-70 contralateral side (isokinetic test)
3. No change in KT scores (+2 or less)
4. Minimal effusion
5. No patellofemoral complaints
6. Satisfactory clinical exam

Goals: Protect healing grafts
Protect patellofemoral joint articular cartilage
Normalize lower extremity strength
Enhance muscular power and endurance
Improve neuromuscular control

- Exercises:
 - Continue previous exercises
 - Emphasis on eccentric quadriceps strengthening
 - Continue closed kinetic chain mini-squats, step-ups, step-downs, lateral lunges, leg press

- Continue knee extension 90 - 40°
 - Hip abduction & adduction
 - Initiate front lunges
 - Calf raises (gastroc and soleus strengthening)
 - Bicycle and stairmaster for endurance
 - Initiate pool running (side shuffle, backward, forward)
 - Initiate walking program
 - Initiate isokinetic exercise 100 - 40° (120-240 °/s spectrum)
- KT-2000 test at week 12
 - Isokinetic testing at week 12 (180° and 300°/s)

V. LIGHT ACTIVITY PHASE (Month 4 – 6)

Criteria to enter Phase V:

1. AROM > 125 degrees
2. Quadriceps strength 70% of contralateral side; flexion/extension ratio 70-79%
3. No change in KT scores (+2 or less)
4. Minimal joint effusion
5. Satisfactory clinical exam

Goals: Enhancement of strength, power, and endurance
Initiate functional and/or sport-specific activity
Prepare for return to functional activities

- Exercises:
 - Continue strengthening exercises – emphasize quadriceps & co-contraction
 - Initiate plyometric program month 4-5
 - Initiate running program month 4-6
 - Initiate agility drills month 5-6
 - Initiate sport-specific training and drills month 5-6
- } If appropriate
- Isokinetic strength test at week 16 & week 18

Criteria to initiate running program:

- Acute reconstruction may begin at 4 – 5 months
- Chronic reconstruction may begin at 5 – 6 months
 1. Satisfactory clinical exam
 2. Unchanged KT test
 3. Satisfactory isokinetic test
 - Quadriceps bilateral comparison (80% or greater)
 - Hamstring bilateral comparison (110% or greater)
 - Quadriceps torque/body weight ratio (55% or greater)
 - Hamstrings/Quadriceps ratio (70% or greater)

4. Proprioception testing 100% of contralateral side
5. Functional hop test > 75% of contralateral leg

VI. RETURN TO ACTIVITY PHASE (Month 6 – 9)

Criteria to return to activities:

1. Satisfactory clinical exam
2. Unchanged KT test
3. Satisfactory isokinetic test
4. Proprioception testing 100% of contralateral side
5. Functional hop test > 80% of contralateral leg

Goals: Gradual return to full-unrestricted sports
Achieve maximal strength and endurance
Normalize neuromuscular control
Progress skill training

- Exercises:
 - Continue strengthening programs
 - Continue proprioception & neuromuscular control drills
 - Continue plyometric program
 - Continue running and agility program
 - Progress sport specific training and drills