

# **Company Overview**


Chanelle Pharma is Irelands largest family-owned indigenous manufacturer of human and animal pharmaceuticals and is Irelands largest wholesaler of veterinary products. Chanelle Pharma strives to bring you an excellent service by providing next day delivery and high-class product support. We continue to reinvest in our product offering, growing significantly year-on-year.

At Chanelle we continuously provide excellent service and support to our customers. If you have any queries please do not hesitate to contact Rory Campbell or your region's Key Account Manager.



**Rory Campbell** - National Sales Manager +353 86 796 8385 rcampbell@chanellegroup.ie







**Key Account** 



**Ross King** 



**Martin Duffy** 



**Marie McGlynn** – North East +353 86 0846214 mmcglynn@chanellegroup.ie

# **Contents**

#### THIS IS A CLICKABLE PDF

CLICK TO JUMP TO YOUR CHOSEN PRODUCT

# **Large Animal Range**

Anthelmintics	5
Anticoccidials	11
Antibiotics	12
Sedatives	13
Non-Steroidal Anti-Inflammatory (NSAID)	15
Supplements	15
Additional Anthelmintics	16
Euthanasia	18
Equine Range	
Anthelmintics	20
Anaesthetics and Sedatives	20
Non-Steroidal Anti-Inflammatory (NSAID)	22
Euthanasia	24
Companion Animal Range	
Anaesthetics and Sedatives	26
Anti-Epileptic Drug	28
Antibiotics	28
Non-Steroidal Anti-Inflammatory	29
Osteoarthritis Multi Modal Treatment	31
Ectoparasiticides & Anthelmintics	
Euthanasia	33
Equipment	
Gloves	35
Dosing and Injectable Applicators	36
Feeding Tubes	37
Syringes and Needles	38

# LARGE ANIMAL RANGE



#### **ANTHELMINTICS**

# Albex® 2.5%

Active Albendazole 2.5% w/v

**Species** Cattle & Sheep

**Description** Is a broad spectrum multi-purpose oral anthelmintic suspension.

 $Treats \& controls \ mature \& \ developing \ immature \ forms \ of gastro-intestinal \ roundworms, lungworms, tapeworms \ and \ adult \ liver \ fluke. Ovicidal \ against \ fluke \ and \ roundworm \ eggs.$ 

Meat Withdrawal Period Cattle: 14 days

Sheep: 4 days

Milk Withdrawal Period Cattle: 60 hours

**Dosage** Cattle: Worms: 30ml/100kg. Fluke & worms: 40ml/100kg

Sheep: Worms: 2ml/10kg. Fluke & worms: 3ml/10kg

Pack Sizes 1 litre, 2.5 litre

**Legal Category** LM (POM from January 2022)



# Albex® 10%

Active Albendazole 10% w/v

**Species** Cattle & Sheep

**Description** Is a broad spectrum multi-purpose oral anthelmintic suspension.

Treats & controls mature & developing immature forms of gastro-intestinal roundworms, lungworms, tapeworms & adult liver fluke. Ovicidal against fluke & roundworm eggs.

Meat Withdrawal Period Cattle: 14 days

Sheep: 4 days

**Dosage** Cattle: Worms: 7.5ml/100kg. Fluke & worms: 10ml/100kg

Sheep: Worms: 0.5ml/10kg. Fluke & worms: 0.75ml/10kg

Pack Sizes 1 litre, 2.5 litre, 5 litre

**Legal Category** LM (POM from January 2022)



# **NEW Albex® Gold**

Milk Withdrawal Period

Active Albendazole 20% w/v

**Species** Cattle

**Description** Treats & controls mature & developing immature forms of

gastro-intestinal roundworms, lungworms, tapeworms & adult liver fluke. Ovicidal against fluke & roundworm eggs

Meat Withdrawal Period Cattle: 7 days

**Dosage** Cattle: Worms: 3.75ml/100kg. Fluke & worms: 5ml/100kg

Cattle: 84 hours

**Pack Sizes** 1 litre, 2.5 litre, 5 litre

**Legal Category** LM (POM from January 2022)



Milk Withdrawal Period

# **ANTHELMINTICS**

# **Animec® Super**

**Active** Ivermectin 1.0% w/v

Clorsulon 10% w/v

**Species** Cattle

**Description** Is a broad-spectrum injectable endectocide. Treats mixed

infestations of adult liver fluke, gastro-intestinal roundworms,

lungworms, eyeworms, mites & lice.

Meat Withdrawal Period 66 days

**Milk Withdrawal Period** Do not use in animals producing milk for human consumption.

**Dosage** 1ml/50kg

**Pack Sizes** 50 ml, 250 ml, 500 ml

**Legal Category** LM (POM from January 2022)



# Animec<sup>®</sup> Injection

**Species** Cattle & Pigs

**Description** Cattle: Treats & controls gastro-intestinal roundworms, lungworms,

eye worms, warbles, mange mites & lice. Pigs: Treats & controls gastro-intestinal roundworms, lungworms, mange mites & lice.

**Meat Withdrawal Period** Cattle: 49 days

Pigs: 28 days

Milk Withdrawal Period Do not use in animals producing milk for human consumption

or in dairy heifers within 60 days prior to calving.

**Dosage** Cattle: 1ml/50kg

Pigs: 1ml/33kg

Pack Sizes 250 ml, 500 ml

**Legal Category** LM (POM from January 2022)



# Animec® Pour-on

Active Ivermectin 0.5% w/v

**Species** Cattle

**Description** Treats & controls gastro-intestinal roundworms, lungworms,

eye worms, warbles, mange mites & lice.

Meat Withdrawal Period Cattle: 28 days

 Milk Withdrawal Period
 Do not use in animals producing milk for human consumption

or in dairy heifers within 60 days prior to calving.

**Dosage** 10ml/100kg

Pack Sizes 1 litre, 2.5 litre, 5 litre

**Legal Category** LM (POM from January 2022)



# ANTHELMINTICS

# Animec® Oral

Active lvermectin 0.08% w/v

**Species** Sheep

**Description** Treats & controls roundworms, lungworms, stomach

worms, mites & nasal bots.

Meat Withdrawal Period

Milk Withdrawal Period Do not use in lactating sheep producing milk for human

consumption. Sheep must not be treated within 60 days prior to the commencement of lactation, if milk is to be used for

human consumption

2.5ml/10kg Dosage **Pack Sizes** 1 litre, 2.5 litre

**Legal Category** LM (POM from January 2022)



# **Bovex**®

Active Oxfendazole 2.265% w/v

Cattle & Sheep **Species** 

**Description** A broad spectrum worm drench for cattle and sheep for the control

of control of mature and immature gastro-intestinal roundworms,

lungworms & tapeworms. This product is also ovicidal against nematode eggs.

**Meat Withdrawal Period** Cattle: 14 days

Sheep: 21 days

Milk Withdrawal Period Cattle: 84 hours

Sheep: Do not use in animals producing milk for human consumption.

Cattle: 20ml/100kg **Dosage** 

Sheep: 2ml/9kg

**Pack Sizes** 1 litre, 2.5 litre, 5 litre, 10 litre

**Legal Category** LM (POM from January 2022)



# Chanaverm Plus®

Levamisole HCL 1.5% w/v Active

Cobalt Sulphate 0.38% w/v

Cattle & Sheep **Species** 

**Description** Oral solution used for the treatment and control of gastro-intestinal  $\,$ 

and pulmonary nematode infections in cattle and sheep.

**Meat Withdrawal Period** Cattle: 18 days

Sheep: 18 days

Milk Withdrawal Period Do not use in animals producing milk for human consumption.

Cattle: 50ml/100kg Dosage Sheep: 5ml/10kg

**Pack Sizes** 5 litre, 10 litre

**Legal Category** LM (POM from January 2022)



# ANTHELMINTICS

# Chan Broad Spec®

**Active** Rafoxanide 2.25% w/v & Levamisole HCL 1.5% w/v

**Species** Cattle & Sheep

**Description** Oral suspension used for the treatment and control of roundworms

> and lungworms. Highly effective against adult (mature) fluke in cattle and sheep and 83% effective against 4-week old immature

fluke in sheep.

**Meat Withdrawal Period** Cattle: 60 days

Sheep: 60 days

Milk Withdrawal Period Do not use in animals producing milk for human consumption.

Dosage Cattle: 50ml/100kg Sheep: 5ml/10kg

**Pack Sizes** 1 litre, 2.5 litre, 5 litre

**Legal Category** LM (POM from January 2022)



# Doramax® Pour-On

Doramectin 0.5%w/v **Active** 

Cattle **Species** 

**Description** For treatment of infestations of gastroint estinal roundworms,

lungworms, eyeworms, warbles, sucking and biting lice,

mange mites and hornfly in cattle.

**Meat Withdrawal Period** 

Milk Withdrawal Period Do not use in pregnant cows or heifers, which are intended

to produce milk for human consumption, within 2 months

of expected parturition.

10ml/100kg Dosage

**Pack Sizes** 1 litre, 3 litre, 5 litre

LM (POM from January 2022) **Legal Category** 



# **Epromec**®

**Active** Eprinomectin 0.5% w/v

**Species** Cattle

**Description** Pour on solution used to treat roundworms, lungworms, warbles,

mange mites, lice and horn flies in beef and dairy cattle.

**Meat Withdrawal Period** 15 days Milk Withdrawal Period Zero hours Dosage 10ml/100kg

**Pack Sizes** 1 litre, 3 litre, 5 litre

LM (POM from January 2022) **Legal Category** 





#### **ANTHELMINTICS**

# Moxodex® Oral

Moxidectin 0.1% w/v **Active** 

Sheep **Species** 

**Description** Oral solution which treats and prevents roundworms

and lungworms in sheep.

**Meat Withdrawal Period** 14 days Milk Withdrawal Period 5 days 2ml/10kg Dosage

**Pack Sizes** 1 litre, 2.5 litre, 5 litre

LM (POM from January 2022) **Legal Category** 



# Ridafluke®

**Species** 

Rafoxanide 3% w/v **Active** 

**Description** Oral suspension which treats & controls fluke infections. Highly

effective against mature fluke in cattle and sheep and 83% effective

against 4-week old immature fluke in sheep.

**Meat Withdrawal Period** Cattle: 60 days

Sheep: 60 days

Cattle & Sheep

Milk Withdrawal Period This product is not permitted for use in animals producing milk

for human consumption, including heifers in the last trimester

of pregnancy.

Cattle: 15ml/40kg **Dosage** 

Sheep: 3ml/8kg

**Pack Sizes** 2.5 litre, 5 litre

**Legal Category** LM (POM from January 2022)



#### Rumenil®

**Active** Oxyclozanide 3.4% w/v

**Species** Cattle

Oral suspension which treats and controls adult liver fluke **Description** 

and tapeworm in cattle.

**Meat Withdrawal Period** 13 Days

Milk Withdrawal Period 108 hours (4.5 days) 3ml/10kg up to 350kg. Dosage

>350kg = 105ml dose

**Pack Sizes** 5 litre

LM (POM from January 2022) **Legal Category** 



#### **ANTHELMINTICS**

# Tribex<sup>®</sup> 5%

Active Triclabendazole 5% w/v

**Species** Sheep

 $\textbf{Description} \hspace{1cm} \textbf{Oral suspension which treats acute, subacute \& chronic fascioliasis}$ 

caused by early immature, immature and adult stages of liver fluke.

Meat Withdrawal Period 56 days

Milk Withdrawal Period Not authorised for use in ewes producing milk for human

consumption including during the dry period. Do not use within 1 year prior to the first lambing in ewes intended to produce milk for

human consumption.

**Dosage** 2ml/10kg

Pack Sizes 800 ml, 2.2 litre, 5 litre

**Legal Category** LM (POM from January 2022)



# Tribex® 10%

Active Triclabendazole 10% w/v

**Species** Cattle

**Description** Oral suspension which treats acute, subacute & chronic fascioliasis

caused by early immature, immature and adult stages of liver fluke.

Meat Withdrawal Period 56 days

 Milk Withdrawal Period
 This product is not permitted for use during lactation in animals

producing milk for human consumption. Not intended for use within 41 days of calving in non-lactating cattle, milk for human consumption may only be taken from 84 hours after calving. If calving occurs before 41 days after treatment, milk for human consumption may only be taken after 41 days plus 84 hours

after the treatment.

Dosage 12ml/100kg

Pack Sizes 800 ml, 2.2 litre, 5 litre

**Legal Category** LM (POM from January 2022)



# Zerofen® 2.5%

Active Fenbendazole 2.5% w/v

**Species** Cattle & Sheep

**Description** Oral suspension which controls mature & developing immature

forms of roundworm & lungworms.

Meat Withdrawal Period Cattle: 14 days

Sheep: 21 days

Milk Withdrawal Period Cattle 96 hour

 $Sheep: do \ not \ use \ in \ sheep \ producing \ milk \ for \ human \ consumption.$ 

Dosage Cattle: 30ml/100kg Sheep: 2ml/10kg

Pack Sizes 1 litre, 2.5 litre

**Legal Category** LM (POM from January 2022)



#### **ANTHELMINTICS**

# Zerofen® 10%

Active Fenbendazole 10% w/v

**Species** Cattle

**Description** Oral suspension which controls mature & developing immature

 $forms\ of\ roundworm\ \&\ lungworms.$ 

Meat Withdrawal Period14 daysMilk Withdrawal Period96 hoursDosage10ml/130kg

Pack Sizes 1 litre, 2.5 litre, 5 litre

**Legal Category** LM (POM from January 2022)



#### **ANTICOCCIDIALS**

#### Chanox Multi®

Active Toltrazuril 5% w/v

**Species** Cattle, Pigs & Sheep

**Description** Oral suspension for the prevention of clinical

signs of coccidiosis.

Meat Withdrawal Period Piglets: 77 days

Calves: 63 days Lambs: 42 days

 Milk Withdrawal Period
 Not permitted for use in lactating animals

producing milk for human consumption.

**Dosage** Calves: 3ml/10kg

Lambs: 4ml/10kg Piglets: 0.4ml/kg

**Pack Sizes** 250 ml, 1 litre

**Legal Category** LM (POM from January 2022)



# **Dycoxan**®

Active Diclazuril 0.25% w/v

**Species** Cattle & Sheep

**Description** Oral suspension for the prevention of clinical signs of

coccidiosis and reduction of oocyst shedding.

**Meat Withdrawal Period** 0 days

Milk Withdrawal Period Not applicable

Dosage Calves: 4ml/10kg

Lambs: 4ml/10kg

Pack Sizes 1 litre, 2.5 litre

**Legal Category** LM (POM from January 2022)



# **ANTIBIOTICS**

# Fenoflox® 5%

Active Enrofloxacin 50 mg/ml

Species Cattle, Pigs, Dogs & Cats

**Description** Treatment of respiratory & alimentary tract infections in

Cattle and Pigs.

**Meat Withdrawal Period** Cattle intravenous: 5 days

Cattle subcutaneous: 12 days

Pigs:13 days

Milk Withdrawal Period Do not use in animals producing milk for human consumption.

Pack Sizes 100 m
Legal Category POM



MA MA

# Fenoflox® 10%

Active Enrofloxacin 100 mg/ml

**Species** Cattle & Pigs

**Description** Treatment of respiratory & alimentary tract infections. Used

for the treatment of local signs associated with pre-acute/acute  $\,$ 

mastitis in dairy cattle.

Meat Withdrawal Period Cattle subcutaneous: 12 days

Cattle intravenous: 5 days Pigs subcutaneous; 13 days

Milk Withdrawal Period Cattle subcutaneous 4 days

Cattle intravenous 3 days

Pack Sizes100 mlLegal CategoryPOM



# Linspec<sup>®</sup>

**Active** Lincomycin

**Species** Cattle (pre-ruminating Calves), Pigs, Dogs & Cats

**Description** Treatment of respiratory infections, intestinal infections, urinary

tract infections, skin infections (including wounds and abscess) and arthritis caused by organisms sensitive to lincomycin and / or

spectinomycin.

**Meat Withdrawal Period** Cattle: 21 days

Pigs: 14 days

Pack Sizes 100ml
Legal Category POM



TO MAN

#### **ANTIBIOTICS**

# Milbotyl<sup>®</sup>

**Active** Tilmicosin 300 mg/ml, Propylene glycol 250 mg/ml

**Species** 

Description For the treatment of pneumonia in cattle and

sheep, ovine mastitis and bovine foot rot.

**Meat Withdrawal Period** Cattle: 70 days

Sheep: 42 days

Milk Withdrawal Period Cattle 36 days

Sheep 18 days

**Pack Sizes** 50 ml VPO **Legal Category** 



# **SEDATIVES**

# Sulpha No 2 Powder®

Sulphadimidine Active

**Species** Calves

**Description** Sulpha No. 2 Powder for oral solution is indicated for use in calves

as an anti-bacterial agent effective against sulfonamide sensitive

organisims.

**Meat Withdrawal Period** 28 days

**Pack Sizes** 200 x 25g sachet

**Legal Category** POM



# Chanazine® 2%

Active Xylazine 20 mg/ml

**Species** Cattle, Horses, Dogs & Cats

**Description** Sedative with analgesic and muscle relaxant properties.

**Meat Withdrawal Period** Cattle:14 days.

Not to be used in horses intended

for human consumption.

Milk Withdrawal Period Do not use in animals producing milk

for human consumption.

**Pack Sizes** 25 ml VPO **Legal Category** 



#### **SEDATIVES**

# Chanazine® 10%

**Active** Xylazine 100 mg/ml

**Species** 

Description Sedative with analgesic and muscle relaxant properties.

The product is for use in horses only, in cases where

sedation is required.

**Meat Withdrawal Period** Not to be used in horses intended

for human consumption.

Milk Withdrawal Period Do not use in animals producing milk

for human consumption.

**Pack Sizes** 50 ml VPO **Legal Category** 



# **Ketamidor®**

Active Ketamine

**Species** Cats, Cattle, Dogs, Horses & Pigs

**Description**  $Ketamine\ is\ a\ dissociative\ anaesthetic.\ Its\ antagonistic\ action\ on$ 

the NMDA-receptors inhibits the neurotransmitter glutamate. The glutamate coupled acetylcholine release blocks the excitability of the neurons and therefore the transmission of neural impulses.

**Meat Withdrawal Period** Zero days Milk Withdrawal Period Zero days

**Legal Category** V/PO



MARAN

# **Medesedan®**

**Pack Sizes** 

**Active** Detomidine hydrochloride 10 mg/ml

**Species** 

**Description** Medesedan injection is an alpha2-drenorecptor agonist, sedative,

analgesic & is used as a pre-anaesthetic. Licensed for IV and IM use.

**Meat Withdrawal Period** Cattle: 2 days

Horses: 2 days

10 ml, 50ml

Milk Withdrawal Period Cattle 12 hours

**Pack Sizes** 5 ml, 20 ml

**Legal Category** VPO



# **NON-STEROIDAL ANTI-INFLAMMATORY (NSAID)**

# **Rifen®**

Active Ketoprofen 100 mg/ml

Species Cattle, Horse & Pigs

**Description** Rifen is a painkiller for large animals. Ketoprofen intervenes in the

synthesis of inflammation mediators by inhibiting the enzyme cyclooxygenase and to some degree also lipoxygenase. As a consequence, no prostaglandins and leukotrienes can be synthesised

which will lead to quick recovery from inflammation.

Meat Withdrawal Period 4 days

Milk Withdrawal Period Zero hours

Pack Sizes 100 ml
Legal Category POM



# Rheumocam<sup>®</sup> Injection



**Description** Used in horses for the relief of equine colic, inflammation

& pain in both acute and chronic musculoskeletal disorders. Used in cattle for acute respiratory infection & acute mastitis. Used for diarrhoea in combination with oral rehydration therapy in calves of over one week of age & young, non-lactating cattle. Used in pigs to reduce the symptoms of lameness & inflammation in non-infectious locomotor disorders. Treats puerperal septicaemia, toxaemia and mastitis-metritis-agalactia syndrome.

**Meat Withdrawal Period** Cattle 15 Days Horses, Pigs 5 days

Milk Withdrawal PeriodCattle 5 DaysPack Sizes100 ml, 250 ml

**Legal Category** POM



# **SUPPLEMENTS**

# **Spasmium® Injection**

Active Metamizole sodium 500 mg, Hyoscine butylbromide 4 mg

**Species** Horses, Cattle, Pigs & Dogs

**Description** Spasmium is for the treatment of spasms or sustained increased

tonus of smooth muscles of the gastrointestinal tract or the urine

and bile excretory organs associated with pain.

Meat Withdrawal Period Horse, Cattle (IV): 12 days Pig (IM): 15 days

Milk Withdrawal Period Cattle (IV) 4 days

Pack Sizes 100 ml
Legal Category POM



#### **SUPPLEMENTS**

# **Cobalt B12 & Selenium**

**Active** Cobalt Acetate 1% w/v

Sodium Selenite 0.025% w/v Vitamin B12 200mcg/ml

**Species** Cattle & Sheep

**Description** Oral supplement, for cattle, calves, sheep and lambs. Provides

essential cobalt, selenium and vitamin B12.

Meat Withdrawal Period0 daysMilk Withdrawal Period0 days

DosageAdult cattle: 20mlSheep: 5ml,Calves: 5mlLambs: 2.5ml

Pack Sizes 1 litre, 2.5 litre

**Legal Category** LM



# **Chanatol**®

Active Cobalt sulphate heptahydrate 0.099% w/v

Propylene glycol to 100% w/v

**Species** Cattle, Sheep & Goats

**Description** Oral solution for the treatment of acetonaemia in cattle

and pregnancy toxaemia in sheep and goats.

Meat Withdrawal Period3 daysMilk Withdrawal Period3 days

**Dosage** Sheep and Goats: 100ml daily for 6 days.

Cattle: 200ml morning and evening for the first day followed

by 100ml twice a day for 3 days.

**Pack Sizes** 1 litre, 2.5 litre, 5 litre

**Legal Category** N/A



# ADDITIONAL ANTHELMINTICS

# **Topimec® Pour-on**

Active Ivermectin 0.5% w/v

**Species** Cattle

**Description** Treats & controls gastro-intestinal roundworms, lungworms,

eye worms, warbles, mange mites & lice.

Meat Withdrawal Period Cattle: 28 days

Milk Withdrawal Period Do not use in animals producing milk for human consumption

or in dairy heifers within 60 days prior to calving.

Dosage 10ml/100kg
Pack Sizes 2.5 litre, 5 litre

**Legal Category** LM (POM from January 2022)



# ADDITIONAL ANTHELMINTICS

# Topimec® Super



**Active** Ivermectin 1.0% w/v

Clorsulon 10% w/v

**Species** Cattle

**Description** A broad-spectrum injectable endectocide. Treats mixed

infestations of adult liver fluke, gastro-intestinal roundworms,

lungworms, eyeworms, mites & lice.

**Meat Withdrawal Period** 66 days

Milk Withdrawal Period Do not use in animals producing milk for human consumption.

Dosage 1ml/50kg

250 ml, 500 ml **Pack Sizes** 

**Legal Category** LM (POM from January 2022)



# **Zeromectin®**



Eprinomectin 0.5% w/v Active

Cattle **Species** 

**Description** Pour on solution used to treat roundworms, lungworms, warbles,

mange mites, lice and horn flies in beef and dairy cattle.

**Meat Withdrawal Period** 15 days Milk Withdrawal Period Zero hours Dosage 10ml/100kg **Pack Sizes** 2.5 litre, 5 litre

**Legal Category** LM (POM from January 2022)



# Triclaben®



**Active** Triclabendazole 10% w/v

**Species** Cattle

**Description** Oral suspension which treats acute, subacute & chronic fascioliasis

caused by early immature, immature and adult stages of liver fluke.

**Meat Withdrawal Period** 56 days

Milk Withdrawal Period This product is not permitted for use during lactation in animals

producing milk for human consumption. Not intended for use within 41 days of calving in non-lactating cattle, milk for human consumption may only be taken from 84 hours after calving. If calving occurs before 41 days after treatment, milk for human consumption may only be

taken after 41 days plus 84 hours after the treatment.

12ml/100kg **Dosage Pack Sizes** 2.2 litre, 5 litre

**Legal Category** LM (POM from January 2022)



# **EUTHANASIA**

# **Euthoxin® Injection**

Active Pentobarbital Sodium 500 mg

**Species** Horses, Ponies, Cattle, Swine, Dogs, Cats, Minks, Polecats, Hares,

Rabbits, Guinea Pigs, Hamsters, Rats, Mice, Chicken, Pigeons, Birds,

Snakes, Tortoises, Lizards, Frogs

**Description** Euthanasia.

**Meat Withdrawal Period** Do not use in animals intended for human or

animal consumption.

Milk Withdrawal Period Do not use in animals intended for human or

animal consumption.

Pack Sizes100 mlLegal CategoryVPO



MARAN+

# **EQUINE RANGE**



# **ANTHELMINTICS**

# **Animec® Oral Paste**

Active Ivermectin 18.7 mg/g

**Species** Horses

**Description** Treats & controls roundworms bloodworms,

pinworms, neck threadworms and bots.

**Pack Sizes** Display tray of 20's

**Legal Category** LM



#### **ANAESTHETICS AND SEDATIVES**

# Iso-Vet®

Active Isoflurane 1000 mg/g

**Species** Horses, Dogs, Cats, Small Mammals,

Ornamental Birds and Reptiles

**Description** Induces and maintain anaesthesia in all types

of veterinary surgery.

Pack Sizes 250 ml

**Legal Category** VPO



# **Butador**®

Active Butorphanol 10 mg/ml

**Species** Horse, Dogs & Cats

**Description** Sedative, analgesic and pre-anaesthetic for horses, can be

combined with xylazine, detomidine and romifidine. Sedative, analgesic and pre-anaesthetic for dogs and cats, can be combined

with medetomidine and ketamine.

Pack Sizes 10 ml, 5 x 10 ml

**Legal Category** POM





# ANAESTHETICS AND SEDATIVES

# Chanazine® 2%

Active Xylazine base 20mg

**Species** Cattle, Horses, Dogs & Cats

**Description** Sedative with analgesic and muscle relaxant properties.

**Meat Withdrawal Period** Cattle 14 days. Not to be used in horses intended for

human consumption.

Milk Withdrawal Period Do not use in animals producing milk

for human consumption.

Pack Sizes 25 ml Legal Category VPO



MATAN

# Chanazine® 10%

Active Xylazine Base 100mg/ml

**Species** Horses

**Description** Sedative with analgesic and muscle relaxant properties. The product

is for use in horses only, in cases where sedation is required.

Meat Withdrawal Period Not to be used in horses intended for

human consumption.

 Milk Withdrawal Period
 Do not use in animals producing milk for

human consumption.

Pack Sizes 50 ml Legal Category VPO



# **Medesedan®**

Active Detomidine hydrochloride 10 mg/ml

(equivalent to 8.36mg detomidine)

**Species** Horse, Cattle

**Description** Medesedan injection is an alpha2-drenorecptor agonist, sedative,

analgesic & is used as a pre-anaesthetic. Licensed for IV and IM use.

Meat Withdrawal Period Cattle: 2 days

Horses: 2 days

Milk Withdrawal Period Cattle 12 hours

**Pack Sizes** 5 ml, 20 ml

**Legal Category** VPO



# **NON-STEROIDAL ANTI-INFLAMMATORY (NSAID)**

# **Ketamidor**®



**Active** Ketamine

**Species** Horse, Cattle, Pig, Dog, Cat

**Description** Ketamine is a dissociative anaesthetic. Its antagonistic action on

the NMDA-receptors inhibits the neurotransmitter glutamate. The glutamate coupled acetylcholine release blocks the excitability of the neurons and therefore the transmission of neural impulses.

Meat Withdrawal PeriodZero daysMilk Withdrawal PeriodZero days

Pack Sizes 10 ml, 50ml

**Legal Category** VPO



# Rheumocam® Granules



**Species** Horses

**Description** Meloxicam granules are for the alleviation of inflamation

and pain in horses.

Pack Sizes 100 x 5g sachet

**Legal Category** POM



# Rheumocam® Oral Suspension



**Species** Horses

**Description** Used for the relief of equine colic & inflammation and pain in

both acute and chronic musculo-skeletal disorders.

Pack Sizes 100 ml, 250 ml

**Legal Category** VPO



# NON-STEROIDAL ANTI-INFLAMMATORY (NSAID)

# Rheumocam® Injection

Meloxicam 20 mg/ml **Active** 

**Species** Cattle, Horses & Pigs

Used in horses for the relief of equine colic, inflammation **Description** 

& pain in both acute and chronic musculoskeletal disorders. Used in cattle for acute respiratory infection & acute mastitis. Used for diarrhoea in combination with oral rehydration therapy in calves of over one week of age & young, nonlactating cattle. Used in pigs to reduce the symptoms of lameness & inflammation in non-infectious locomotor disorders. Treats puerperal septicaemia, toxaemia and

mastitis-metritis-agalactia syndrome.

Meat Withdrawal Period Cattle 15 Days

Horses, Pigs 5 days

Milk Withdrawal Period Cattle 5 Days

**Pack Sizes** 100 ml, 250 ml

**Legal Category** POM



# Rifen® Injection

**Active** Ketoprofen 100 mg/ml **Species** Horse, Cattle & Swine

Description Rifen is a painkiller for large animals. Ketoprofen intervenes

in the synthesis of inflammation mediators by inhibiting the enzyme cyclooxygenase and to some degree also lipoxygenase. As a consequence, no prostaglandins and leukotrienes can be synthesised which will lead to quick recovery from inflammation.

**Meat Withdrawal Period** 4 days

Zero hours Milk Withdrawal Period

100 ml **Pack Sizes** 

**Legal Category** POM





# Spasmium<sup>®</sup> Injection

Active Metamizole Sodium Monohydrate 500mg

Hyoscine butylbromide 4 mg

**Species** Horses, Cattle, Pigs & Dogs

**Description**  $Spasmium\,is\,for\,the\,treatment\,of\,spasms\,or\,sustained\,increased$ 

tonus of smooth muscles of the gastrointestinal tract or the

urine and bile excretory organs associated with pain.

**Meat Withdrawal Period** Horse, cattle (IV): 12 days

Pig (IM): 15 days

Milk Withdrawal Period Cattle (IV): 4 days

**Pack Sizes** 100 ml **Legal Category** POM



# **EUTHANASIA**

# **Euthoxin® Injection**

Active Pentobarbital Sodium 500 mg

**Species** Horses, Ponies, Cattle, Swine, Dogs, Cats, Minks, Polecats, Hares,

Rabbits, Guinea Pigs, Hamsters, Rats, Mice, Chicken, Pigeons, Birds,

Snakes, Tortoises, Lizards and Frogs

**Description** Euthanasia

Meat Withdrawal Period Do not use in animals intended for human or

animal consumption.

Milk Withdrawal Period Do not use in animals intended for human or

animal consumption.

Pack Sizes 100 ml Legal Category VPO



MARKET +

# COMPANION ANIMAL RANGE



# **ANAESTHETICS AND SEDATIVES**

# Iso-Vet®

Active Isoflurane 1000 mg/g

Species Horses, Dogs, Cats, Small Mammals,

Ornamental Birds and Reptiles

**Description** Induces and maintain anaesthesia in all types

of veterinary surgery.

Pack Sizes250 mlLegal CategoryVPO



# Bupaq<sup>®</sup>

Active Buprenorphine 0.3 mg

**Species** Dogs & Cats

**Description** Bupaq is recommended for pre-and postsurgical analgesia in long

lasting interventions with expected moderate to severe pain.

Pack Sizes 10 ml Legal Category VPO



# **Butador**®

**Species** 

Active Butorphanol 10 mg

**Description** Sedative, analgesic and pre-anaesthetic for horses, can be

Horses, Dogs, & Cats

combined with xylazine, detomidine and romifidine. Sedative, analgesic and pre-anaesthetic for dogs and cats, can be combined

with medetomidine and ketamine.

Pack Sizes 10 ml, 5 x 10 ml

**Legal Category** POM



# **ANAESTHETICS AND SEDATIVES**

# Chanazine® 2%

**Active** Xylazine base 20mg

**Species** Cattle, Horses, Dogs & Cats

**Description** Sedative with analgesic and muscle relaxant properties.

**Meat Withdrawal Period** Cattle 14 days. Not to be used in horses intended for

human consumption.

Milk Withdrawal Period Do not use in animals producing milk

for human consumption.

Pack Sizes 25 ml Legal Category VPO



MAMA

# **Medetor®**

Active Medetomidine hydrochloride 1 mg

**Species** Dogs & Cats

**Description** Alpha2-drenorecptor agonist, it is a sedative, analgesic and

is used as a pre-anaesthetic.

Pack Sizes 10 ml Legal Category VPO



# **Insistor**®

**Active** Methadone Hydrochloride 10mg

(equivalent to 8.9mg Methadone)

**Species** Dogs & Cats

**Description** Insistor is indicated for analgesia as a premedicant for general

anaesthsia or for neuroleptanalgesia in combination with a

neuroleptic drug.

Pack Sizes 10 ml

**Legal Category** VPO



#### **ANAESTHETICS AND SEDATIVES**

# **Revertor®**

MI

Active Atipamezole hydrochloride 5 mg

**Species** Dogs & Cats

**Description** Alpha2-antagonist, it is used for the reversal of the sedative

effects of medetomidine (Medetor®) and dexmedetomidine.

in dogs & cats.

Pack Sizes 10 ml
Legal Category POM



# **ANTI-EPILEPTIC DRUG**

# **Epityl**®

**Active** Phenobarbital 60 mg

**Species** Dogs

**Description** Phenobarbital is an antiepileptic agent for the prevention of seizures

due to generalized epilepsy in dogs.

Pack Sizes 100 tabs, 500 tabs

**Legal Category** POM



# **ANTIBIOTICS**

# Fenoflox® 5%

**Active** 

Enrofloxacin 50 mg/ml

**Species** Cattle, Pigs, Dogs & Cats

**Description** Treatment of respiratory & alimentary tract infections in Cattle

and Pigs. Treats bacterial infections of the alimentary, respiratory and urogenital tracts, skin, wound infections and otitis externa in

dogs & cats.

**Meat Withdrawal Period** Cattle intravenous: 5 days

Cattle subcutaneous: 12 days

Pigs: 13 days

 Milk Withdrawal Period
 Do not use in animals producing milk

for human consumption.

Pack Sizes 100 ml Legal Category POM



A A

#### **ANTIBIOTICS**

# **Linspec**®

**Active** Lincomycin (as lincomycin hydrochloride) 50mg

Spectinomycin (as spectinomycin sulfate tetrahydrate) 100mg

 $\textbf{Species} \hspace{1cm} \textbf{Cattle (pre-ruminating Calves), Pigs, Dogs \& Cats}$ 

**Description** Treatment of respiratory infections, intestinal

infections, urinary tract infections, urinary infections, skin infections (including wounds and abscess) and arthritis caused by organisms

sensitive to lincomycin and/ or spectinomycin.

Meat Withdrawal Period Cattle: 21 days

Pigs: 14 days

Pack Sizes 100 ml

**Legal Category** POM



A MA

# **NON-STEROIDAL ANTI-INFLAMMATORY (NSAID)**

# Canidryl® Snap Tab

**Active** Carprofen

**Species** Dogs

**Description** Canidryl flavoured tablets are used for the reduction of

inflammation and pain caused by musculoskeletal disorders and degenerative joint disease and as a follow up to parenteral analgesia in the management of post-operative

pain following soft tissue surgery in dogs.

**Strength** 20 mg, 100 mg

Pack Sizes 100 tabs
Legal Category POM



# Rheumocam® Chewable Tablets

Active Meloxicam 2.5 mg

**Species** Dogs

**Description** Used to alleviate inflammation and pain in both acute and chronic

musculo-skeletal disorders and the management of post-

operative pain. following soft tissue surgery in dogs.

Strength1 mg, 2.5 mgPack Sizes100 tabsLegal CategoryPOM



# **NON-STEROIDAL ANTI-INFLAMMATORY (NSAID)**

# Rheumocam<sup>®</sup> Injection

Active Meloxicam 5 mg/ml

**Species** Dogs & Cats

**Description** For the alleviation of inflammation and pain in both

acute and chronic musculo-skeletal disorders in dogs.

Pack Sizes 10 ml, 20 m

**Legal Category** POM



# Rheumocam® Oral Suspension for Dogs

Active Meloxicam 1.5 mg/ml

**Species** Dogs

**Description** Alleviation of inflammation and pain in both acute and

chronic musculo-skeletal disorders in dogs.

**Pack Sizes** 15 ml, 42 ml, 100 ml, 200 ml

**Legal Category** POM



# **Rheumocam® Oral Suspension for Cats**

Active Meloxicam 0.5mg/ml

**Species** Cats

**Description** Description To relieve the pain and inflammation associated

with chronic musculoskeletal conditions and acute injury

Pack Sizes 5ml, 15ml

**Legal Category** POM



# **NON-STEROIDAL ANTI-INFLAMMATORY (NSAID)**

# **Spasmium® Injection**



Hyoscine butylbromide 4 mg

**Species** Horses, Cattle, Pigs, Dogs

**Description** Spasmium is for the treatment of spasms or sustained increased

tonus of smooth muscles of the gastrointestinal tract or the

urine and bile excretory organs associated with pain.

**Meat Withdrawal Period** Horse, cattle (IV): 12 days Pig (IM): 15 days

Milk Withdrawal Period Cattle (IV): 4 days

Pack Sizes 100 ml
Legal Category POM



# **OSTEOARTHRITIS MULTI MODAL TREATMENT**

# Osteopen®

Active Pentosan Polysulphate Sodium 100mg/ml

**Species** Dogs

**Description** For the treatment of lameness and pain of degenerative joint disease/

osteoarthritis (non-infectious arthrosis) in the skeletally mature dogs.

**Strength** 100 mg/ ml

Pack Sizes 10ml
Legal Category POM



# **ECTOPARASITICIDES & ANTHELMINTICS**

#### Fleatix<sup>®</sup>

**Active** Fipronil

**Species** Dogs & Cats

**Description** Fleatix is an easy to use spot-on solution for the treatment of

flea and tick infestations. Fleatix Spot-On Solution has a persistent insecticidal efficacy for up to 5 weeks in fleas and can be used as part

of a treatment strategy for Flea Allergic Dermatitis, where a veterinary surgeon has diagnosed this.

**Strength** Small Dog, Medium Dog, Large Dog, Extra Large Dog & Cat

Pack Sizes 10 x 3's
Legal Category LM



# **ECTOPARASITICIDES & ANTHELMINTICS**

# Fleatix® Combo

Active Fipronil (S)-methoprene

**Species** Dogs & Cats

**Description** Easy to use spot on solution that kills fleas, ticks, biting lice and stops

the development of flea eggs, larvae and pupae within the home.

 $\textbf{Strength} \hspace{1cm} \textbf{Small Dog, Medium Dog, Large Dog, Extra Large Dog \& Cat} \\$ 

Pack Sizes 10 × 3's
Legal Category LM



# **NEW Chanhold®**

**Active** Selamectin

**Species** Dogs & Cats

 $\textbf{Description} \hspace{1cm} \textbf{Treats and prevents infections of heartworms, fleas, ear mites,} \\$ 

sarcoptic mange, and certain types of ticks in dogs, and prevents heartworms, fleas, ear mites, hookworms, and roundworms in cats.

**Strength** Cats and Dogs less than 2.5 kg, Toy Dogs, Small Dogs, Medium Dogs,

Large Dogs. Medium Cat and Large Cat.

Pack Sizes 10 x 3's
Legal Category POM



# **NEW Imoxat®**

Active Imidacloprid & Moxidectin

**Species** Dogs, Cats & Ferrets

**Description** Treats and prevents a wide range of both internal and external

parasites

**Strength** Small Dogs, Medium Dogs, Large Dogs, Extra large dogs,

Small Cat and Large Cat.

Pack Sizes 10 x 3's
Legal Category POM



# **ECTOPARASITICIDES & ANTHELMINTICS**

# Prazitel® Plus

**Active** Praziquantel 50 mg/Pyrantel 144 mg/Febantel 150 mg

**Species** 

Description Flavoured tablets which treats roundworms, hookworms,

whipworms, tapeworms and infection in dogs and puppies.

**Pack Sizes** 24's, 100's, 200's

**Legal Category** LM



# Prazitel® Plus XL



Active Praziquantel 175 mg/Pyrantel 504 mg/Febantel 150 mg

**Species** Dogs

**Description** Flavoured tablets which treat roundworms, hookworms,

whipworms and tapeworm infection in dogs and puppies.

**Pack Sizes** 10's, 50's

**Legal Category** LM



# Prazitel® Cat



Praziquantel 20 mg/Pyrantel 230 mg3 Active

**Species** 

**Description** Treats roundworms and tapeworm infections in cats.

24's, 104's

**Pack Sizes** LM

**Legal Category** 



# **EUTHANASIA**

# **Euthoxin® Injection**

Active Pentobarbital Sodium 500 mg

**Species** Horses, Ponies, Cattle, Swine, Dogs, Cats, Minks, Polecats, Hares,

Rabbits, Guinea Pigs, Hamsters, Rats, Mice, Chicken, Pigeons, Birds,

Snakes, Tortoises, Lizards, and Frogs

**Description** Euthanasia.

**Meat Withdrawal Period** Do not use in animals intended for human or animal consumption.

Milk Withdrawal Period Do not use in animals intended for human or animal consumption.

Pack Sizes 100 ml Legal Category VPO



MARKET +

# **EQUIPMENT**

# **Equipment**

# **GLOVES**

# **Albex Milking Gloves**

Pack Size	Size	Product Code
100	Medium	XX1624
	Large	XX1623
	X Large	XX1625

#### Description

Powder and latex free. Ambidextrous. Exceptional comfort and strength. Ideal for milking, hoof trimming and general farm use. Black in



# **Chanelle Disposable Gloves 100's**

Pack Size	Size	Product Code
100	Arm Length	XX0038

#### **Description**

Single use. Ambidextrous. Non-sterile. Extra-sensitive. Suitable for calving, lambing and examinations.



# **Kruse Latex Gloves - Powder Free**

Pack Size	Size	<b>Product Code</b>
100	Small	KR1472
	Medium	KR1475
	Large	KR1787

#### Description

Smooth, durable, elastic latex examination gloves with high sensitivity. Rolled cuff. Non-sterile. Ambidextrous.



# **Krutex Latex Gloves**

Pack Size	Size	Product Code
100	Small	KR1428
	Medium	KR1409
	Large	KR1408
	X Large	KR1069

#### Description

 $High \, quality, latex \, examination \, gloves. \, Smooth, \, elastic \, and \, with \, high \,$ sensitivity. Rolled cuff, non-sterile, ambidextrous. Cornstarch powder. Thickness at the fingers: 0.16 mm. Thickness at the palm: 0.15 mm. AQL:  $\,$ 



# **DOSING & INJECTABLE APPLICATORS**

# Pour On Gun 60ml

Species	Size	Product Code
Cattle	60ml	XX1612

#### Description

Adjustable volume and easy calibration. Complete with chemical resistant tubing. Impact-resistant, durable hard-plastic casing. Comfortable handle design for easy gripping.



# **Chanox Dosing Gun 30 ml**

Species	Size	Product Code
Cattle/Sheep	10ml	XX3678
Cattle/Sheep	30ml	XX3679

#### Description

Top quality, specially designed drenching gun for use when treating calves with Chanox product. Also compatible across full Chanelle range. Strenghtened hard-plastic casing and easy calibration.



# Hookmatic 80 Drencher 80 ml

Species	Size	<b>Product Code</b>
Cattle	80ml	QT1010

#### Description

Reinforced steel hook design for easy use with large cattle. High volume delivery system. Comfort grip incorporated into design of both hook and gun hand pieces. Chemical resistant tubing



# **Chanelle Auto Drencher**

Species	Size	Product Code
Cattle/Sheep	20ml	QT1005
Cattle/Sheep	30ml	QT1006

#### Description

Automatically refills, saving time and labour. Use for both sheep and cattle. Durable hard-plastic design and easily calibrated.



# **DOSING & INJECTABLE APPLICATORS**

# **Dosing Drencher N/Auto**

Species	Size	<b>Product Code</b>
Cattle	150ml	AZ1002
Cattle/Sheep	60ml	AZ1003

#### Description

 $\label{thm:continuous} \mbox{High volume delivery system. Extra strength hard-plastic design. Easy calibration.}$ 



# **Chanelle Auto Injector**

Species	Size	Product Code
Cattle/Sheep	12ml	XX0275

#### Description

Adjustable volume, durable design. Automatically refills, saving time and labour. Can be used in both cattle and sheep. Comfort-grip handle.



# **FEEDING TUBES**

# **Lamb Feeding Tube**

Species	Product Code
Sheep	XX1087

#### Description

Syringe and catheter for feeding weak lambs.



# **Chanelle Calf Feeding Tube & Bag**

Species	Product Code
Cattle	XX1006

#### Description

Stomach tube and feeding bag for feeding colostrum/electrolytes to weak calves. Enlarged, rounded end for animal comfort and ease of use. Stopper valve on tube to control flow.



# **Equipment**

# **SYRINGES & NEEDLES**

# **BD Syringes**

Product	Pack Size	Size	Product Code
- Syr BD 1ml (300013)	100	1ml	MC1034
- Syr BD 2ml Emerald (307727 )	100	2ml	MC1009
- Syr BD 5ml Emerald (307731)	100	5ml	MC1002
- Syr BD 10ml Emerald (307736)	100	10ml	MC1001
- Syr BD Disp 20ml (300613)	120	20ml	MC1012
- Syr BD Disp 30ml (301231)	60	30ml	MC1014
- Syr BD (Inj) 50ml (300866)	60	50ml	MC1007
- Syr BD 2ml with 21G x 5 needle (300325)	100	2ml	MC1033



#### Description

 $Single-use. \ Clear \ cylinder \ with \ a \ easily \ readable \ scale. \ Complies \ with \ international \ performance \ standards. \ CE \ marked.$ 

Our Mission is to provide affordable medicines that extend and improve the lives of people and animals.

